

RECEIVED
CENTRAL FAX CENTER


JUL 24 2009

Appl. No. 10/584877
Reply to Office Action dated 1/26/09**Amendments to the Claims:**

This Listing of Claims will replace all prior versions and listings of claims in the application. No new matter has been added.

Listing of Claims:

1. (Currently Amended) A method for the treatment ~~or prevention~~ of autoimmune arthritis comprising administering to a patient an effective amount of a coumestans compound of formula I or pharmaceutically acceptable salts thereof, or an extract containing the coumestans compound of formula I or pharmaceutically acceptable salts thereof



wherein

R₁ represents H, OH, or methoxyl;

R₂ represents H, OH, or C₁-C₈ alkyl;

R₃ and R₄ are each independently selected from the group consisting of H, halogen, OH, and methoxyl.


2. (Previously Presented) The method of claim 1, wherein the pharmaceutically acceptable salt of the compound of formula I are formed with the acids selected from the group consisting of chlorhydric acid, hydrobromic acid, sulfuric acid, citric acid, tartaric acid, phosphoric acid, lactic acid, pyruvic acid, acetic acid, succinic acid, oxalic acid, fumaric acid, maleic acid, ketosuccinic acid, methane-sulfonic acid, ethyl-sulfonic acid, benzene sulfonic acid, and isethionic acid.

3. (Previously Presented) The method of claim 1, wherein the compound of formula I or the extract is extracted from the plant *Compositae*.

Appl. No. 10/584877
Reply to Office Action dated 1/26/09

4. (Previously Presented) The method of claim 3, wherein the plant *Compositae* is selected from *Eclipta prostrata* Linn, *Wedelia chinensis*, and *Eclipta alba*.

5. (Previously Presented) The method of claim 1, wherein said compound of formula I is wedelolactone as shown in formula II:



II

6. (Previously Presented) The method of claim 1, wherein the arthritis is selected from the group consisting of rheumatic arthritis, rheumatoid arthritis, and osteoarthritis.

7. (Previously Presented) The method of claim 1, wherein the compound of formula I is produced by an extraction method comprising the steps of:

- (a) extracting the fruits, leaves, or branches of the *Compositae* plant with 95±3% ethanol, thereby producing ethanol extract;
- (b) dissolving the above ethanol extract in 5-300 volumes of H₂O at 50-80°C, filtering to remove the precipitates and collecting the H₂O phase;
- (c) extracting the H₂O phase in step (b) with acetic ester and collecting the acetic ester phase;
- (d) concentrating and drying the acetic ester phase in step (c) to produce the precipitates;
- (e) eluting the precipitates in (d) on a silica gel column with a 5:1 to 1:2 gradient of petroleum ether/acetone mixture and collecting the fraction eluted with 1:1 petroleum ether/acetone;
- (f) concentrating the fraction eluted in step (e) to produce concentrated residue;
- (g) eluting the concentrated residue in step (f) on a silica gel column with a 5:1 to 1:2 gradient of dichloromethane/acetone mixture and collecting the fraction eluted with 3:1 dichloromethane/acetone;

Appl. No. 10/584877
Reply to Office Action dated 1/26/09

(h) eluting the fraction eluted in step (g) on a silica gel column with a 20:10:1 to 5:10:1 gradient of toluene-acetone-formate mixture and collecting the fraction eluted with 10:10:1 toluene-acetone-formate;

(i) eluting the fraction eluted in step (h) on a silica gel column with a 30:1 to 1:1 gradient of dichloromethane/methanol mixture and collecting the fraction eluted with 20:1 dichloromethane/methanol; and

(j) recrystallizing the fraction eluted in step (i) with ethanol, thereby producing the coumestans compounds of formula I as precipitates.

8. (Original and Withdrawn) A method for producing coumestans, comprising the steps of:

(a) extracting the fruits, leaves, or branches of the *Compositae* plant with 95±3% ethanol, thereby producing ethanol extract;

(b) dissolving the above ethanol extract in 5-300 volumes of H₂O at 50-80°C, filtering to remove the precipitates and collecting the H₂O phase;

(c) extracting the H₂O phase in step (b) with acetic ester and collecting the acetic ester phase;

(d) concentrating and drying the acetic ester phase in step (c) to produce the precipitates;

(e) eluting the precipitates in (d) on a silica gel column with a 5:1 to 1:2 gradient of petroleum ether/acetone mixture and collecting the fraction eluted with 1:1 petroleum ether/acetone;

(f) concentrating the fraction eluted in step (e) to produce concentrated residue;


(g) eluting the concentrated residue in step (f) on a silica gel column with a 5:1 to 1:2 gradient of dichloromethane/acetone mixture and collecting the fraction eluted with 3:1 dichloromethane/acetone;

(h) eluting the fraction eluted in step (g) on a silica gel column with a 20:10:1 to 5:10:1 gradient of toluene-acetone-formate mixture and collecting the fraction eluted with 10:10:1 toluene-acetone-formate;

Appl. No. 10/584877
Reply to Office Action dated 1/26/09

(i) eluting the fraction eluted in step (h) on a silica gel column with a 30:1 to 1:1 gradient of dichloromethane/methanol mixture and collecting the fraction eluted with 20:1 dichloromethane/methanol; and

(j) recrystallizing the fraction eluted in step (i) with ethanol, thereby producing the coumestans compounds of formula I as precipitates



I


wherein,

R₁ represents H, OH, or methoxyl;

R₂ represents H, OH, or C₁-C₈ alkyl;

R₃ and R₄ are each independently selected from the group consisting of H, halogen, OH, and methoxyl.

9. (Original and Withdrawn) A dietary supplement comprising 0.05-50wt% of the compound of formula I, or the pharmaceutically acceptable salts thereof:



I

wherein

R₁ represents H, OH, or methoxyl;

R₂ represents H, OH, or C₁-C₈ alkyl;


R₃ and R₄ are each independently selected from the group consisting of H, halogen, OH, and methoxyl;

and a bromatologically acceptable carrier.

Appl. No. 10/584877
Reply to Office Action dated 1/26/09

10. (Original and Withdrawn) A pharmaceutical composition for treating arthritis, comprising:

(a) 0.05-90wt% compound of formula I or pharmaceutically acceptable salts thereof, or extracts containing the coumestans compound of formula I or pharmaceutically acceptable salts thereof, as the main active ingredient,



wherein

R₁ represents H, OH, or methoxyl; and

R₂ represents H, OH, or C₁-C₈ alkyl;

R₃ and R₄ are each independently selected from the group consisting of H, halogen, OH, and methoxyl; and

(b) one or more active ingredients selected from the group consisting of: acetaminophen, diclofenac, ibuprofen, indomethacin, meloxicam, ketoprofen, sulindac, auranofin, naproxen, nabumetone, piroxicam, meclofenamic acid, chlofenamic acid, mefenamic acid, pirofen, fenbufen, tolmetin, flufenamide acid, methocarbamol, nimesulide, celecoxib, rofecoxib, aceclofenac methotrexate, gold salts, salazosulfadimidine, penicillamine, chloroquine, tripterygium wilfordii, ciclosporin, cyclophosphamide, and glucocorticoids; and

(c) a pharmaceutically acceptable carrier.